

DOI: 10.34854/ICPAF.51.2024.1.1.254

KINETIC PROCESSES INVOLVING ELECTRONICALLY EXCITED MOLECULAR NITROGEN IN THE MIDDLE ATMOSPHERES OF TITAN AND EARTH ^{*)}

Kirillov A.S.

Polar Geophysical Institute, Apatity, Russia, kirillov@pgia.ru

Molecular nitrogen N₂ is the main molecular gas in the atmospheres of the Earth, Titan (a moon of Saturn), Triton (a moon of Neptune) and Pluto. In the Earth's atmosphere, the second gas in total concentration is molecular oxygen O₂; in the atmospheres of the other mentioned planets it is methane CH₄.

A study was carried out of the kinetics of the triplet A³Σ_u⁺, B³Π_g, W³Δ_u, B³Σ_u⁻, C³Π_u states of molecular nitrogen at altitudes of the middle atmosphere of Titan of 50-250 km during the precipitation of cosmic rays into the atmosphere. The calculations take into account intramolecular and intermolecular electron energy transfer during inelastic collisions of electronically excited molecular nitrogen with N₂, CH₄ and CO molecules. The interaction constants of electronically excited molecular N₂(A³Σ_u⁺) with N₂ and CO molecules are calculated according to quantum chemical approximations and show good agreement with the available experimental data [1]. The interaction of electronically excited N₂ molecules with molecules of methane CH₄, acetylene C₂H₂, ethylene C₂H₄, ethane C₂H₆ in the middle atmosphere of Titan at altitudes of 50-250 km was studied. The dominance of reactions with metastable molecular nitrogen N₂(A³Σ_u⁺) in the formation of C₂H and C₂H₃ radicals at these altitudes was shown for the first time [2].

Similar kinetic calculations involving triplet electron-excited molecular nitrogen were carried out for the Earth's middle atmosphere of 30-80 km during the precipitation of high-energy relativistic electrons into the atmosphere [3]. The interaction constants of metastable molecular nitrogen N₂(A³Σ_u⁺) with oxygen molecules O₂ were calculated and compared with the available experimental data [4]. The emission intensities of the bands of the first positive and second positive N₂ systems during the precipitation of high-energy electrons were calculated. It is shown that there is a significant decrease in the emission intensities of the bands of the first positive system with decreasing altitude due to the influence of collision processes on the populations of vibrational levels of the N₂(B³Π_g) molecule. The influence of intermolecular processes of energy transfer from N₂(A³Σ_u⁺) on the formation of singlet oxygen and the emission of the Atmospheric and Infrared atmospheric bands of O₂ at altitudes of the Earth's middle atmosphere was studied.

References

- [1]. Kirillov A.S. Intermolecular electron energy transfer processes in the collisions of N₂(A³Σ_u⁺, v=0-10) with CO and N₂ molecules. *Chemical Physics Letters*, 2016, v.643, p.131–136.
- [2]. Kirillov A.S., Werner R., Guineva V. Studying the electron kinetics of molecular nitrogen in the middle atmosphere of Titan during the precipitation of cosmic rays. *Bulletin of the Russian Academy of Sciences: Physics*, 2023, v.87, p.985-993.
- [3]. Kirillov A.S., Belakhovsky V.B. The kinetics of N₂ triplet electronic states in the upper and middle atmosphere during relativistic electron precipitation. *Geophysical Research Letters*, 2019, v.46, p.7734-7743.
- [4]. Kirillov A.S., Belakhovsky V.B. The kinetics of O₂ singlet electronic states in the upper and middle atmosphere during energetic electron precipitation. *Journal of Geophysical Research: Atmosphere*, 2021, v.126, e2020JD033177.

^{*)} [abstracts of this report in Russian](#)